



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search ☒ for

Limits

Preview/Index

History

Clipboard

Details

Display ☒ Show ☒ Sort by ☒ Send to ☒All: 1 Review: 0 ☒

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: Diabetologia. 2005 Jun;48(6):1207-15. Epub 2005 Apr 30.Related Articles,
Links**Alterations of lipids and apolipoprotein CIII in very low density lipoprotein subspecies in type 2 diabetes.****Hiukka A, Fruchart-Najib J, Leinonen E, Hilden H, Fruchart JC, Taskinen MR.**

Department of Medicine, Division of Cardiology, Helsinki University Hospital and Biomedicum, Haartmaninkatu 8, 00029, Helsinki, Finland.

AIMS/HYPOTHESIS: Very low density lipoprotein (VLDL) particles are heterogeneous, comprising two main subspecies, VLDL 1 (Sf 60-400) and VLDL 2 (Sf 20-60). The aim of the study was to examine the distribution and composition of VLDL subspecies in type 2 diabetes. **SUBJECTS, MATERIALS AND METHODS:** We studied the composition and concentration of triglyceride-rich lipoproteins (TRLs) in 217 type 2 diabetic patients and 93 control subjects between 50 and 75 years of age. Lipoprotein subspecies were separated by density-gradient ultracentrifugation. Apolipoprotein (apo) CIII and apo E in plasma and apo CIII in TRL subspecies were measured by nephelometry and apo CII in serum by a commercial kit using a single radial immunodiffusion method. **RESULTS:** The concentrations of VLDL 1, VLDL 2 and intermediate density lipoprotein were significantly increased in type 2 diabetes subjects, the change being most marked for VLDL 1. There was a strong linear correlation between VLDL 1 triglycerides and plasma triglycerides in both groups ($r = 0.879$, $p < 0.001$ and $r = 0.899$, $p < 0.001$). Diabetic subjects had markedly higher plasma ratios of apo CII:apo CIII and apo CIII:apo E. Despite elevated plasma apo CIII, type 2 diabetic subjects had a relative deficiency of apo CIII in all TRL subspecies, suggesting profound disturbances of apo CIII metabolism. **CONCLUSIONS/INTERPRETATION:** The elevation of VLDL 1 triglycerides is the major determinant of plasma triglyceride concentration in normal subjects and in type 2 diabetic individuals. Both apo CIII and apo E metabolism are disturbed in type 2 diabetes.

PMID: 15864534 [PubMed - indexed for MEDLINE]

Display ☐ Show ☐ Sort by ☐ Send to ☐

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Oct 18 2005 10:52:14